

Investigating and concentrating iron-chromium-nickel ores of the Malkinsk deposit and of manganese ores of the Labinsk deposit in Northern Caucasus. A. A. Glazkovskiy. *Izvestiya Akademiya Nauk SSSR, Tekhnicheskaya Mekhanika* (Engl. Mech. Engineering), 1932, 170-61. The Malkinsk deposit consists of magnetite, hematite and brown bog iron. It contains: Fe, 41.01-41.29; Ni, 0.20-0.30; Cr, 0.06-0.11; Mn, 2.58-3.21; Al₂O₃, 14.1-15.76; CaO, 1.03-1.18; MgO, 2.44-2.56; MnO, 0.05-0.08; Cr₂O₃, 1.12-1.16; TiO₂, 0.06-0.09; P₂O₅, 0.01-0.04 and loss on heat treating 7.41-11.15. It is possible to prep. Fe concentrates by flotation and magnetic sepn.; Ni is left in the tailings. The approx. compn. of the Labinsk ore is SiO₂, 44.35-55.22; Fe, 6.12-10.57; Mn, 12.32-15.77; MnO₂, 17.22-26.11; S, 0.02-0.11; P, 0.05-0.11 and H₂O, 4.56-7.22%. The ore can be used to produce silicomanganese or ferro-manganese.

A. A. Reichtung

a

8

Copper-containing magnetite deposits of the Urals and
their exploitation V. A. Glukhovskiy Soviet Inst. 9.
No. 6, 66-1011930 The deposits contain 70% Fe.
Cu can be obtained as a by-product B. H. M.

AS & SLA METALLURGICAL LITERATURE CLASSIFICATION

U.S. PATENT OFFICE

RESEARCH AND DEVELOPMENT

RESEARCH AND DEVELOPMENT

ALADINSKIY, P.I.; ARONSKIND, S.Sh.; GLAZKOVSKIY, V.A.; KVASKOV, A.P.;
SUVOPOV, F.S.; SHMANENKOV, I.V., redaktor; BASHANOV, V.A.,
redaktor; SERGEYEVA, N.A., redaktor; MANINA, M.P., tekhnicheskii
redaktor

[Results of the organization and work of an ore-dressing laboratory]
Opyt organizatsii i raboty obogatitel'noi laboratorii. Trudy lab.
geol.upr. no.3:3-57 1952. [Microfilm] (MLRA 7:11)
(Ore dressing)

GLAZKOVSKIY, V.A.; ROZHKOVA, Ye.V., redaktor; RAZUMNAYA, Ye.G., redaktor;
BABINTSEV, N.I., redaktor; POPOV, N.D., tekhnicheskii redaktor.

[Geological and mineralogical basis for metallurgical assaying
of iron ores from deposits of different genesis] Geologo-minera-
logicheskie osnovy tekhnologicheskoi otsenki rud mestorozhdenii
zheleza razlichnogo genezisa. Moskva, Gos. nauchno-tekhn. izd-vo
lit-ry po geologii i okhrane neдр, 1954. 181 p. (MLRA 7:12)
(Iron ores)

GLAZKOVSKIY, YE.V.

Controlling the cleanness of spinneret orifices by means of an
opillascop. Khim.volok. no.1:50-51 '66. (NIRA 13:6)

1. Kalininskiy filial Vsesoyuznogo nauchno-issledovatel'skogo
institut iskusstvennogo volokna.
(Rayon spinning)

GORYACHKO, G.V.; LARIONOV, N.I.; GLAZKOVSKIY, Yu.V.

Ultrasonic cleaning of spinnerets. Khim.volok. no.1:51-52
'60. (MIRA 13:6)

1. Kalininskiy pedinstitut (for Goryachko, Larionov). 2. Kalininskiy
filial Vsesoyuznogo nauchno-issledovatel'skogo instituta iskusst-
vennogo volokna (for Glazkovskiy).
(Rayon spinning)
(Ultrasonic waves--Industrial applications)

MIKHAYLOVA, G.S.; GLAZKOVSKIY, Yu.V.; GRAFOV, V.V.

Internal dyeing of cuprammonium fiber using ultrasonic dispersion
of pigments. Khim.volok. no.2:61-62 '62. (MIRA 15:4)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut steklyanogo
volokna.

(Dyes and dyeing--Rayon)

ADDOV, F.L.; ZIBUNYANOV, F.V.; GLA KOVCHIK, Yu.V.

Using the chemical and spectral analysis methods in studying
the process of alkali saponification of polyacrylonitrile.
Dokl. Akad. Nauk SSSR, 1965, no. 4, 81-87, 165.
(MIRA 18:9)

1. Markovskiy khimicheskiy institut.

GLAZNAY, Edward, gen., inz.

Two new construction concepts for filters. Izvestiya 17 no.12:
647-652 '61.

S/081/62/000/025/045/120
B166/B101

AUTHOR: Glazmak, Edward

TITLE: Two new continuous filter designs

PERIODICAL: Referativnyy zhurnal. Khimiya, no. 23, 1962, 587, abstract
23176 (Przegl. chem., v. 17, no. 12, 1961, 647-652 [Pol.])

TEXT: The multi-section vacuum filter is a modified drum vacuum filter in which the fixed, suspension-filled trough is absent. The filter is designed in the form of a horizontal drum, having its outer surface divided into segments by longitudinal partitions. The segments are closed at the ends, forming trough-shaped cells on the drum's surface. Inside these cells are located the filtering elements which consist of two perforated plates representing two sides of an isosceles triangle placed with its base on the bottom of the trough-shaped cell. The suspension to be separated drops from above into each cell in succession when the vacuum filter drum is rotated on its axis; from the cell the filtrate is sucked inside the drum. The operating cycle of this filter is the same as that of an ordinary drum vacuum filter. The horizontal continuous filter is a plate rotating about its

Card 1/2

3/04/61/000/025/045/120
3166/8101

Two new continuous filter designs

vertical axis. Vertical cylinders with perforated walls are mounted around the circumference of the plate and, inside these cylinders the filter mesh is placed. When the plate rotates the lower holes in the cylinders join up in succession with the pipelines along which the suspension and wash waters are arriving. The filter cake is removed from the cylinder by means of a Z-shaped knife fixed at the center of each cylinder which is actuated by meshing with a gear wheel. [Abstracter's note: Complete translation.]

Card 2/2

WILLIAM L. B. ...

...

...

...

...

GLAZMAN, B.A.; SAVINYKH, A.G.; GLADKOVA, A.A.; LYUKHANOV, G.F.; KUNDIN, V.M.;
MERTINS, I.P.

Automation of hydrolysis processes. Gidroliz. i lesokhim. prom.
17 no.7:26-28 '64. (SIRA 17:11)

1. Krasnodarskiy gidroliznyy zavod (for Glazman, Savinykh, Gladkova,
Lyukhanov). 2. Proyektno-konstruktorskoye byuro Severo-Kavkazskogo
soveta narodnogo khozyaystva (for Kundin, Mertins).

a L 9785-66

ACC NR: AP5028541

SOURCE CODE: UR/0286/65/COO/020/0151/0151

AUTHORS: Kavalero¹¹¹v, A. A.; Mirosh¹¹¹nenko, P. A.; Nor¹¹¹inskiy, Ye. Ya.; Sidorov, K. I.; Glazman, B. M.; Kry¹¹¹mchanskiy, F. G.; Ivanov, I. I.

ORG: none

TITLE: Earth digging machine for ditch digging. Class 8h, No. 175895 [announced by Special Construction Bureau No. 1 of the State Committee on Construction, Road Building and Municipal Machinery Construction at GOSSTROYe of the SSSR (Osoboye konstruktorskoye byuro No. 1 gosudarstvennogo komiteta stroitel'nogo, dorozhnogo i kommunal'nogo mashinostroyeniya pri GOSSTROYe SSSR)]

SOURCE: Byulleten' izobreteniy i tovarnykh znakov, no. 20, 1965, 151

TOPIC TAGS: earth handling equipment, construction equipment, tractor, transportation equipment

ABSTRACT: This Author Certificate presents a ditch digging machine. The machine includes a tractor and a supporting frame on which are mounted a cutter, a discharge cone, a thrower with rotating mantle, a plow-type wideners, and a drive (see Fig.1). To decrease the metal and power requirements, the digger is con-

Card 1/2

UDC: 621.879.48.867.9

L 9785-66

ACC NR: AP5028541




Fig. 1. 1 - Tractor; 2 - lifting frame;
3 - face cutter; 4 - discharge cone;
5 - thrower; 6 - rotating thrower mantle;
7 - plow-shaped wideners; 8 - drive;
9 - movable cutting blades; 10 - mantle
support; 11 - levers of face cutter.

structured with a face cutter on the hub of which movable cutting blades are mounted. These are automatically rotated when the face cutter rotation is reversed. The cutter has a common drive with the thrower whose rotating mantle is mounted on a central support. A second feature has the rotation mechanism for the movable blades executed in the form of a pneumatic cylinder which is mounted in the sleeve of the lifting frame and which acts on levers rigidly connected to the blades of the face cutter. Orig. art. has: 1 figure.

SUB CODE: 13/

SUBM DATE: 09Jul64

Card 2/2

GLAZMAN, B.S.; YELISHEV, A.P.

Precision casting of 3Kh2B8 steel. Lit. proizv. no. 6:44-45 Je '61.
(MIRA 14:6)

(Precision casting)

GLAZMAN, B.S.

Factors having an effect on the quality of brass castings made
under pressure. Lit. proizv. no.8:33-34 Ag '62, (MIRA 15:11)
(Die casting) (Brass founding)

Radiotekhnika, 11, Fasc. 7, 1962

642

117

calculated the nominal power of the tube in this case. It is only 1.5 mW. The increase of selectivity of the selective circuit of the input circuit of the transmitter means a considerable improvement of the resulting energy coefficient. The selective circuit of the end stage of the transmitter with increased selectivity is then located. The purpose of the selectivity of this circuit system by a simple means is to ensure that immediately tuned circuits in the whole circuit of the transmitter. It is a great advantage to construct the circuit system in form of a band filter with the largest peaks at the transmission boundaries of the band. The circuit diagram of such a filter as well as its equivalence scheme are discussed on the basis of a drawing. Finally, the equivalent resistance of the tube which is caused by this circuit system, is determined. A further advantage offered by the selective circuit is the protection of the output of the television transmitter against the frequency of the sound emitter in the case of the operation of a sound antenna. Experimental verification resulted in a sufficiently accurate agreement with theoretically computed relations. The selective circuitry does not have a noticeable broadening of the transmission band of the end stage without any reduction of its power. The here discussed scheme was realized in many television stations of the Soviet Union.

INSTITUTION:

GLAZMAN, E. S.

E. S. GLAZMAN: "Absorber of the higher order for television transmitters." Scientific Journal devoted to "Radio Eng.", No. 120, Transmitter 1-10, Moscow, 9 Sep. 56

Analysis of the peculiarities of the operating regions of resonators in an echo-absorber circuit. The analysis carried out shows that an echo-absorber reduces the wave reflection level by 20 - 25 times at the central frequency and by 5 - 7 times at the edges of the working frequency band in the most unfavorable case (for operation in one television channel). This permits a considerable simplification in the construction and regulation of all antenna-feeder apparatus in VLF television radio stations.

GLAZMAN, E.S.

Feeder-echo absorber for television transmitters. Radiotekhnika 14
no.2:3-16 F '59. (MIRA 12:1)
(Television--Transmitters and transmission)

Source: Mathematical Reviews,

Continued

the von
s about
e Chap-
sets of
al and
chapter
stons is
for self
a final
mental
on the
xorems
tegrals
sected
twists
djoint
rather
spec-
The
perce
imple
case
sical
quar-
cent
strong
multip
these
stems
in a
tion
sult
ous)
the
are
are
fers
that
the
ides
the
any
for

GLAZMAN, I. M.

Glazman, I. M. On the spectrum of linear differential
operators. Doklady Akad. Nauk SSSR (N.S.) 80: 133
1981

... B. ...

Mathematical Reviews
Vol. 10, No. 11
Nov. 1963
Analysis

✓ Glazman, I. M. On the character of the spectrum of one-dimensional singular boundary problems. Doklady Akad. Nauk SSSR (N.S.) 87, 5-8 (1952). (Russian)

Let $St(A)$ be the spectrum of an operator A and let $D_d(A)$ be the discrete part and $C_d(A)$ the continuous part of the spectrum, $\sigma = D_d(A) \cup C_d(A)$. Let L be a differential operator on $(0, \infty)$ and L^* a corresponding self-adjoint operator. The author proves a number of theorems. I. If $q(x) \rightarrow 0$ as $x \rightarrow \infty$, then $St(L)$, with $L = (D^2 + q(x)^2)^{1/2}$, contains $\lambda > 0$. If $\limsup_{x \rightarrow \infty} q(x) = \liminf_{x \rightarrow \infty} q(x)$, for $x \rightarrow \infty$, is finite, then for all $\lambda > 0$, $C_d(L) \cap \{\lambda, \lambda + \epsilon\} \neq \emptyset$. II. Let

$$H(x) = \sum_{k=0}^n (-1)^k f_k(x) x^{n-k} y^{(k)}$$

and let $f_k(x) = (-1)^k f_k^{(n)}(x) > 0$ for $k = 0, 1, \dots, n$; $f_k \in C^1$, $k = 0, 1, \dots, n-1$. Then $St(L) \cap [0, \infty) \neq \emptyset$. III. Let II hold and

$$\lim_{x \rightarrow \infty} f_k(x) = 0, \quad k = 0, 1, \dots, n-1.$$

Then $(-\infty, 0) \cap C_d(L) = \emptyset$. Many further results are given. V. I. Lomovskii (Moscow, Moscow U.)

GLAZMAN, I. M.

USSR/Mathematics - Eigenvalues

11 Nov 52

"The Character of the Spectrum of Multidimensional Singular Boundary-Value Problems," I. M. Glazman, Khar'kov Polytech Inst imeni Lenin

"Dokl Ak Nauk SSSR" Vol 87, No 2, pp 171-174

Investigates the disposition of the continuous part of the spectrum of the boundary-value problem connected with the multidimensional differential operation of the second order of the following form. $\Delta u = -\Delta u + q(P)u$, considered in the entire n-dimensional

Euclidean space. Cites allied work of A. M. Molchanov ("Dokl Ak Nauk SSSR" 83, 17 (1952)). Submitted by Acad A. N. Kolmogorov 10 Sep 52.

PA 245T72

245T72

GLAZMAN, I. M.

USSR J MS

Glazman, I. M. On an application of the method of decomposition to multidimensional singular boundary problems. Mat. Sb. N.S. 35(77), 231-246 (1954). (Russian)
The author extends to the multidimensional case the method of decomposition previously considered by the author [Dokl. Akad. Nauk SSSR (N.S.) 30, 153-156 (1951); 87, 5-8 (1952); MR 13, 654; 14, 1088] for the study of the nature of the spectrum for one-dimensional singular boundary-value problems.

The operator $[Lu] = -\Delta u + q(P)u$ considered in a domain Ω is called singular if Ω is unbounded or if $q(P)$ is not con-

tinuous in $\bar{\Omega}$. The case where Ω is bounded by a closed surface σ is considered and the unbounded Ω is denoted by $\Omega(\sigma, \infty)$. Self-adjoint operators L associated with l on $\Omega(\sigma, \infty)$ are considered. The effect on the continuous spectrum is considered for a change in σ . A change in $l=l_1$ to $l=l_2$ by modifying $q=q_1(P)$ to $q_2(P)$, where $q_2(P)-q_1(P) \rightarrow 0$ as $P \rightarrow \infty$ leaves the continuous spectrum of L_1 identical with that of L_2 if the domain of L_2 is that of L_1 . Further results are given.

N. Levinson (Cambridge, Mass.).

X-P/W

GLAZMAN I. M.

USSR/ Mathematics - Spectral functions

Card 1/1 Pub. 22 - 6/62

Authors : Glazman, I. M., and Nayman, P. B.

Title : On the convex cover of orthogonal spectral functions

Periodical : Dok. AN SSSR 102/3, 445 - 448, May 21, 1955

Abstract : Some problems are discussed connected with the construction of a set of all spectral functions of a differential system:

$$-y'' + q(x)y - \lambda y = 0, \quad y'(0) = hy(0) \quad (0 \leq x < \infty)$$

which is considered as a convex set of functions. Definitions of a spectral, and an orthogonal spectral function are given. Seven references: 6 USSR and 1 Swiss (1940-1954).

Institution : V. I. Lenin Polytechnical Institute, Kharkov, and The Aviation Institute, Kharkov.

Presented by: Academician S. L. Sobolev, January 27, 1955

AKHIEZER, N.I., prof. (Khar'kov); GLAZMAN, I.M. (Khar'kov)

Certain classes of continuous functions generating Hermite-
positive kernels. Uch.zap.KhGU 80:205-217 '57.

(MIRA 12:11)

(Functions, Continuous)

ADJACK:

Ginzburg, I.M.

20-2-3/62

ATTN:

An Analogue of the Extension Theory of Derivation Operators and a Non-Symmetric Unidimensional Boundary Value Problem on a Half-Axis.
(Ob odnom analoge teorii rasshireniy kraevykh operatorov i nesimmetricheskoy odnomernoy kraevoy zadache na polosei)

REMARKS:

Uchenye Zapiski SSSR, 1957, Vol.115, Nr 2, pp. 214-216

ABSTRACT:

H signifies here Hilbert's space, J - a certain adjunction operator in H (i.e., an operator defined everywhere in H, which satisfies the conditions $(Jf, Jg) = (g, f)$ and $J^2f = f$ for any f and g from H.) First the author gives several definitions: 1) A linear operator acting in H is called J-symmetric, when $(Af, Jg) = (f, JAf)$ applies to all f and g from the definition domain D_A of the operator A. The J-symmetric operator A with a definition domain dense in H is called J-selfadjoint, when $JAJ^* = A^*$ applies. 3) A linear operator A acting in H is called dissipative, when for any $f \in D_A$ the condition $\text{Im} (Af, f) \geq 0$ applies. In this paper altogether 3 theorems are given: Theorem 1) In order that a dissipative J-symmetric operator with a definition domain dense in H is J-selfadjoint, it is necessary and sufficient that its defect number is equal to zero. Theorem 2) Any J-symmetric operator A with a definition domain dense in H permits an extension to a dissipative operator \tilde{A} self-adjoint in J. Theorem 3) When the condition $\text{Im } p_k(x) \geq 0$ ($k = 0, 1, \dots, n$; $0 \leq x < \infty$)

Card 1/2

GLAZMAN, I. M., Doc Phys-Math Sci--(diss) "Direct Methods of Qualitative Spectrum Analysis of Singular Differential Operators." Khar'kov, 1953. 21 pp, (in Higher Ed Ukr SSR. Khar'kov Order of Labor Red Banner State Univ im A. M. Gor'kiy), 150 copies, Bibliography at the end of the text (25 titles). (KL, 40-58, 112)

AUTHOR: Glazman, I.M.

SOV/42-13-3-7/41

TITLE: On the Developability in Terms of the System of Eigenelements of Dissipative Operators (O razlozhimosti po sisteme sobstvennykh elementov dissipativnykh operatorov)

PERIODICAL: Uspekhi Matematicheskikh Nauk, 1958, Vol 13, Nr 3, pp 179-181 (USSR)

ABSTRACT: The system $\{\varphi_k\}_{k=1}^{\infty}$ of elements of the Hilbert space H is called a base of Riesz for its closed linear closure if for every $f = \sum_{k=1}^n c_k \varphi_k$ there holds the inequality

$$m \sum_{k=1}^n |c_k|^2 \leq \|f\|^2 \leq M \sum_{k=1}^n |c_k|^2,$$

where M and m are positive constants independent of f .

The author improves a result due to Mukminov [Ref 2].

Theorem: Let A be a bounded dissipative operator with the system of eigenelements $\{\varphi_k\}_{k=1}^{\infty}$, $(\varphi_k, \varphi_k) = 1$. Let $\{\lambda_k\}_{k=1}^{\infty}$

be the corresponding sequence of eigenvalues. If

Card 1/2

On the Developability of the System of Eigenelements
of Dissipative Operators

30V/42-13-3-7/41

$$\sum_{j,k=1}^{\infty} \frac{\operatorname{Im} \lambda_j \operatorname{Im} \lambda_k}{|\lambda_j - \bar{\lambda}_k|^2} < \infty \quad (j \neq k),$$

then the system $\{\varphi_k\}_{k=1}^{\infty}$ is a base of Riesz for its closed

linear closure.

There are 3 Soviet references.

SUBMITTED: February 21, 1957

Card 2/2

AUTHOR: GLAZMAN, I.M.

20-3-1/59

TITLE: Oscillation Theorems for Differential Equations of Higher Order and the Spectrum of the Corresponding Differential Operators (Ostsillyatsionnye teoremy dlya differentsial'nykh uravneniy vysshikh poryadkov i spektr sootvetstvuyushchikh differentsial'nykh operatorov)

PERIODICAL: Doklady Akademii Nauk ^{SSSR}, 1958, Vol. 118, Nr. 3, pp. 423-426 (USSR)

ABSTRACT: Given the equation

$$(1) \quad l[y] \equiv \sum_{k=0}^n (-1)^{n-k} [p_k(x) y^{(n-k)}]^{(n-k)} = \lambda y \quad (p_0(x) \neq 0, 0 \leq x < \infty).$$

Lemma: Let \tilde{L} be a selfadjoint operator generated by the operation l and U be the negative part of the spectrum of \tilde{L} . In order that the set U is bounded from below and discrete, it is necessary and sufficient that for every $\varepsilon > 0$ there exists an α such that the quadratic functional

$$\phi_{\varepsilon}[y] \equiv \int_{\alpha}^{\infty} l[y] \bar{y} \, dx + \varepsilon \int_{\alpha}^{\infty} |y|^2 \, dx$$

is non-negative. In order that U is finite, it is necessary and sufficient that for a certain α : $\phi_0[y] \geq 0$.

Card 1/4

Oscillation Theorems for Differential Equations of Higher
Order and the Spectrum of the Corresponding Differential
Operators

20-3-1/59

Definition: (1) is called oscillatory if for every α there exists a solution of (1) which at the right of α has more than one n -tuply zero.

Theorem: In order that (1) is non-oscillatory for $\lambda = \lambda_0$, it is necessary and sufficient that the part of the spectrum of \tilde{L} which lies at the left of $\lambda = \lambda_0$ is an infinite set.

Theorem: For every $\delta > 0$ let $\int_{M_{k\delta}} |p_k^*(x)| dx < \infty$ ($k=1,2,\dots,n$).

Here let $p^*(x)$ be the negative part of the function $p(x)$ and let $M_{k\delta}$ be the set of values of x for which $|p^*(x)| \geq \delta$.

Then for $\lambda < 0$, (1) is non-oscillatory.

Let $p_k(x) = a_k = \text{const}$ and K_a denote the set of points

$Q(a_1, a_2, \dots, a_n)$ of the n -dimensional coefficient space which correspond to the equations (1) being non-oscillatory for $\lambda = 0$. By the transformation $x = \ln t$, $y = x^{1-\lambda} z$ the

Card 2/4

functional $\phi_0[y]$ which corresponds to an equation (1) with

Oscillation Theorems for Differential Equations of Higher
Order and the Spectrum of the Corresponding Differential
Operators

20-3-1/59

constant coefficients is brought to the form

$$\tilde{\Phi}_0[z] = \int_{\alpha'}^{\infty} |z_t^{(n)}|^2 dt + \sum_{k=1}^n \int_{\alpha'}^{\infty} b_k t^{-2k} |z_t^{(n-k)}|^2 dt$$

where b_k are linear functions of the a_k :

$$(2) \quad b_k = \varphi_k(a_1, \dots, a_n) \quad (k=1, 2, \dots, n)$$

Theorem: Let K_b be a convex set arising from K_a by the transformation (2). Let $b_k' = \liminf_{x \rightarrow \infty} p_k(x)$ and $b_k'' = \limsup_{x \rightarrow \infty} p_k(x)$.

If $Q(b_1', b_2', \dots, b_n') \in K_b$, then for $\lambda = 0$, (1) is non-oscillatory.

If $Q(b_1'', b_2'', \dots, b_n'') \notin K_b$, then for $\lambda = 0$, (1) is oscillatory.

Card 3/4 Theorem: For $q(x) \geq -\alpha_n^2 x^{-2n}$ the equation

Oscillation Theorems for Differential Equations of Higher
Order and the Spectrum of the Corresponding Differential
Operator

20-3-1/59

(3) $(-1)^n y^{(2n)} + q(x)y = \lambda y$
is non-oscillatoric and for $q(x) < -(\alpha_n^2 + \delta)x^{-2n}$, $\delta > 0$

it is oscillatoric. Here $\alpha_n = \frac{(2n-1)!!}{2^n}$.

Further four theorems on the equation (3) are given.
6 Soviet and 2 foreign references are quoted.

ASSOCIATION: Kharkov Polytechnical Institute im.V.I.Lenina (Khar'kovskiy
politekhicheskiy institut im.V.I.Lenina)

PRESENTED: By S.N.Bernshteyn, Academician, 15 July 1957

SUBMITTED: 15 July 1957

AVAILABLE: Library of Congress

Card 4/4

AUTHOR: Glazman, I.M. (Kharkov) 20-119-3-5/65

TITLE: On the Negative Part of the Spectrum of Onedimensional and Multidimensional Differential Operators on Vector Functions (Ob otritsatel'noy chasti spektra odnomernykh i mnogomernykh differentsial'nykh operatorov nad vector-funktsiyami)

PERIODICAL: Doklady Akademii Nauk, 1958, Vol 119, Nr 3, pp 421-424 (USSR)

ABSTRACT: The author generalizes his results of [Ref 1]. Let $\vec{L}_2(0, \infty)$ be the Hilbert space of the vector functions $\vec{y}(x) = \{y_k\}_{k=1}^m$ ($m < \infty$) with the scalar product

$$(\vec{y}, \vec{z}) = \int_0^\infty \sum_{k=1}^m y_k(x) \overline{z_k(x)} dx$$

and $l[\vec{y}]$ a differential operation

$$(1) \quad l[\vec{y}] = (-1)^n \vec{y}^{(2n)} + Q(x) \vec{y} \quad (0 \leq x < \infty)$$

where $Q(x)$ is an Hermitian matrix of m -th order. The smallest

Card 1/3

On the Negative Part of the Spectrum of Onedimensional Differential Operators on Vector Functions and Multidimensional

20-119-3-5/65

and the highest eigenvalue of $Q(x)$ are $\mu^*(x)$ and $\nu(x)$ respectively. Let \tilde{L} denote an arbitrary self-adjoint extension of the operator with a minimum definition domain which is generated in $\tilde{L}_2(0, \infty)$ by (1). Let be $f^*(x) =$

$= \min \{0, f(x)\}$.

Theorem: Let M_δ be the set of all x for which $|\mu^*(x)| \geq \delta$

If for each $\delta > 0$ there holds the inequality $\int_{M_\delta} |\mu^*(x)| dx < \infty$

then the negative part of the spectrum of \tilde{L} is bounded from below and discrete.

Theorem: The negative part of the spectrum of \tilde{L} consists of finitely many eigen values, if one of the following conditions is satisfied

a) $\mu(x) \geq -\alpha_n^2 x^{-2n}$ for large x , $\alpha_n = \frac{(2n-1)!}{2^n}$ b) for each $\delta > 0$ it is $\int_{M_\delta} x^{2n-1} |\mu^*(x)| dx < \infty$.

where M_δ is the set of all x , for which $|\mu^*(x)| \geq (\alpha_n^2 - \delta) x^{-2n}$

Card 2/3

On the Negative Part of the Spectrum of Onedimensional and Multidimensional Differential Operators on Vector Functions 20-119-3-5/65

for a certain $p \gg 1$ it holds $\int_0^\infty x^{2np-1} \|\mu^*(x)\|^p dx < \infty$.

Theorem: If for a $\delta > 0$ and a natural r for sufficiently

great x it holds: $\nu(x) < -\frac{1}{4x^2} - \frac{1}{4x^2 \ln^2 x} - \dots - \frac{1+\delta}{4x^2 \ln^2 x \dots \ln_r^2 x}$,

where $\ln_k x = \ln \ln_{k-1} x$, then the negative part of the spectrum of \bar{L} consists of finitely many eigen values.

A third theorem contains conditions under which the negative part of the spectrum of \bar{L} consists of infinitely many values.

The fourth theorem is a generalization of the first one to differential operators of the type $l[\vec{u}] = -\Delta \vec{u} + Q(p) \vec{u}$.

There are 8 references, 5 of which are Soviet, and 3 American.

Card 3/3

ASSOCIATION: Khar'kovskiy politekhnicheskii institut imeni V.I. Lenina
(Kharkov Polytechnical Institute imeni V.I. Lenin)

PRESENTED: October 24, 1957, by S.N. Bernshteyn, Academician

SUBMITTED: October 24, 1957

Subject: [illegible]

[illegible text]

32444

16.3400

S/044/61/000/010/008/051
C111/C222

AUTHORS: Glazman, I.M., and Zhikhor', N.A.

TITLE: The reversion of singular dissipative differential operators
of second order

PERIODICAL: Referativnyy zhurnal. Matematika, no. 10, 1961, 21,
abstract 10 B 100. ("Uch. zap. Khar'kovsk. gos. ped. in-t",
1957, 21, 9-24)

TEXT: In the $L^2(0, \infty)$ the author considers the differential operator L
generated by the differential expression $l(y) = -y'' + q(x)y$
(closure of the operator $L_1 y = l(y)$ given on the finite functions), where
 $q(x)$ is a complex-valued function with a non-negative imaginary part which
is summable on every finite interval $[0, 1]$. It is proved that for
 $\text{Im } \lambda < 0$ for the corresponding equation

$$-y'' + q(x)y = \lambda y \quad (1)$$

the basic results of Weyl remain true, i.e. there occurs the case either
of the boundary point or the boundary circle. In the first case (1) has
only one (up to a constant factor) solution of $L^2(0, \infty)$, in the second
Card 1/2

32444

The reversion of singular dissipative ... S/044/61/000/010/008/051
C111/C222

case all solutions of (1) belong to $L^2(0, \infty)$. Furthermore the author investigates complete dissipative extensions L of the operator L which satisfy the conditions $\tilde{L} \supseteq L$, $\text{Im} (\tilde{L}f, f) \geq 0$

$$f \in D_{\tilde{L}}, (\tilde{L} - \lambda L)D_{\tilde{L}} = L^2(0, \infty).$$

It is stated that in the case of the boundary point all such extensions are given by boundary conditions of the type $y'(0) = hy(0)$, $\text{Im } h \geq 0$, and that the resolvent of an arbitrary such extension is a bounded integral operator with a Carleman kernel; in the case of the boundary circle the resolvent of an arbitrary such extension is a kernel of Hilbert-Schmidt.

[Abstracter's note : Complete translation.]

Card 2/2

GLAZMAN, I. M.; SKACHEK, B. Ya.

On the discrete part of the Laplacian spectrum in regions
of limiting cylindricality. Dokl. AN SSSR 147 no.4:760-763
D '62. (MIRA 16:1)

1. Khar'kovskiy politekhnicheskij institut im. V. I. Lenina.
Predstavleno akademikom S. N. Bernshteynom.

(Operators(Mathematics)) (Eigenvalues)

AM4036551

BOOK EXPLOITATION

S/

Glazman, Izrail' Markovich

Direct methods in qualitative spectrum analysis of singular differential operators (Pryamy*ye metody* kachestvennogo spektral'nogo analiza singulyarny*kh differentsial'ny*kh operatorov), Moscow, Fizmatgiz, 1963, 338 p. illus., biblio., index. 6,500 copies printed.

TOPIC TAGS: qualitative spectral analysis, singular differential operator, mathematics, Schroedinger operator, differential equation

PURPOSE AND COVERAGE: This monograph presents the fundamentals and uses of direct methods of investigating the nature of the spectrum of singular differential operators. It covers journal literature of the past 10-15 years. The Schroedinger operator occupies a central place among the operators that are considered. The book can be of interest not only to the mathematician, but also to the physicist. It is within the reach of students in the advanced courses and graduate students wishing to get acquainted with qualitative spectral theory of differential equations.

TABLE OF CONTENTS [abridged]:

Card 1/2

AM1036551

Foreword -- 7

Introduction -- 9

Ch. I. Method of splitting and general theorems on the spectrum -- 17

Ch. II. Spectrum of unidimensional two-member differential operators -- 130

Ch. III. Spectrum of unidimensional differential operators of a general type -- 197

Ch. IV. Spectrum of multidimensional differential operators -- 221

Ch. V. Spectrum of the Schroedinger differential operator and solutions of equations -- 264

Ch. VI. Some additional problems of qualitative spectral analysis -- 298

Bibliography -- 327

Subject index -- 338

SUB CODE: MA

SUBMITTED: 24Oct63

NR REF SOV: 069

OTHER: 042

DATE ACQ: 16Apr64

Card 2/2

BR

ACCESSION NR: AP4043894

S/0179/64/000/004/0126/0128

AUTHOR: Glazman, I. M. (Khar'kov); Shteynvol't, L. I. (Khar'kov)

TITLE: Eliminating the natural frequencies of a vibrating system in the dangerous resonance zones by variation of its parameters

SOURCE: AN SSSR. Izvestiya. Mekhanika i mashinostroyeniye, no. 4, 1964, 126-128

TOPIC TAGS: vibration, natural frequency, resonance, resonance elimination, vibrating system, programming, rigidity

ABSTRACT: A method is described for eliminating the natural frequencies from the dangerous resonance zone of linearly rotating vibrating systems with a finite number of degrees of freedom. This problem is solved by using digital computers on the basis of the Jacobi-Sylvester theorem (F. R. Gantmakher). S is considered to be a simple or complex rotating vibrating system with a finite number of degrees of freedom. The matrix of the S system is designated as $A=A(u)$ so that the squares of the natural frequency of the S system are the roots of the equation:

Card 1/3

$$\text{Det } (A(u) - \lambda I) = 0$$

(1)

ACCESSION NR : AP4043894

The authors first determine whether the existing rigidity is sufficient. If not, it may then be assumed that the rigidity varies either in one direction or in both directions. Systems of equations are then evolved for these cases as a basis for computer programs. For one direction:

$$D_1(u_1, \dots, u_m), D_2(u_1, \dots, u_m), \dots, D_n(u_1, \dots, u_m) \quad (2)$$

And considering the rigidity to be constant:

$$P_1(u), P(u), \dots, P_n(u) \quad (3)$$

For two directions:

$$\Delta_1(u_1, \dots, u_m), \Delta_2(u_1, \dots, u_m), \dots, \Delta_n(u_1, \dots, u_m) \quad (4)$$

And considering the rigidity to vary:

(5)

$$(\max_u \min_x P(x, u) > 0$$

Card 2/3

ACCESSION NR: AP4043894

"The programming was performed by A. A. Motornaya, E. M. Livshitz and A. I. Kononenko to whom the author wishes to express his thanks". Orig. art. has: 8 equations.

ASSOCIATION: none

SUBMITTED: 10Nov63

ENCL: 00

SUB CODE: ME, DP

NO REF SOV: 002

OTHER: 000

Card 3/3

ACCESSION NR: AP4016497

S/0020/64/154/005/1011/1014

AUTHOR: Glazman, I. M.

TITLE: On gradient relaxation for non-quadratic functionals

SOURCE: AN SSSR. Doklady*, v. 154, no. 5, 1964, 1011-1014

TOPIC TAGS: non-quadratic functional, gradient relaxation, Euclidean space, Hilbert space, descent method

ABSTRACT: The method of descent, proposed by L. V. Kantorovich (DAN 48, no. 7, 1945) was investigated by him and other authors for the case of a positive definite quadratic functional in Euclidean or Hilbert space. Beyond the boundaries of the class of such functionals, the method of descent presents a problem. The present paper is devoted to one particular question of the problem of descent. Its basic aim is the effective construction of a universal algorithm of gradient relaxation for the class K of all functions $\Phi(x)$ of point $x(x_1, x_2, \dots, x_n)$ of Euclidean space E_n , which satisfy the following three conditions: (1) $\Phi(x)$ is twice continuously differentiable; (2) $\Phi(\infty) = \infty$; (3) $\Phi(x)$ has a unique stationary point \bar{x} . The known methods of gradient relaxation for quadratic

Card 1/3

ACCESSION NR: AP4016497

functionals do not carry over to this case. Thus, for example, the method of least descent would require at each step, for the determination of the next factor of complete relaxation, the solution of non-linear equation, which itself requires an infinite computation process. Realization of a stationary procedure of gradient relaxation would require an initial evaluation of an upper bound for the maximum M of the operator norm of the Hessian matrix

$$H(x) = \left(\frac{\partial^2 \Phi}{\partial x_i \partial x_k} \right)_{i,k=1}^p$$

in the region $\Omega \subset \mathbb{E}_p$ of all x satisfying $\Phi(x) < \Phi(x_0)$. An effective construction of an auxiliary algorithm for such an estimate, valid for the whole class K , would again require, it seems, the construction of a minimizing sequence (now for the functional $\|H(x)\|$ and with estimates of the rapidity of convergence, which leads to a vicious circle. In the general case of a non-stationary relaxation process

$$x_{n+1} = x_n - \gamma_n \nabla \Phi(x_n), \quad (n = 0, 1, 2, \dots) \quad (3)$$

Card2/3

ACCESSION NR: AP4016497

which will be called length of the relaxation path the sum of the series

$$\sum_{n=0}^{\infty} |x_{n+1} - x_n|. \quad (4)$$

The algorithm described in this article may be utilized for the construction of minimizing sequences in variation problems. Orig. art. has: 10 equations.

ASSOCIATION: Fiziko-tekhnicheskii institut nizkikh temperatur, akademii nauk USSR (Physico - Engineering Institute of Low Temperature, Academy of Science, USSR).

SUBMITTED: 25Sep63

DATE ACQ: 12Mar64

ENCL: 00

SUB CODE: MA

NO REF SOV: 005

OTHER: 002

Card 3/3

L 55956-65 EWT(d)/T Pg-4/Ph-4 IJP(c)

ACCESSION NR: AP5010818

UR/0020/64/161/004/0750/0752

AUTHOR: Glazman, I. M.

TITLE: Relaxation on surfaces with saddle points

SOURCE: AN SSSR. Doklady, v. 161, no. 4, 1968, 750-752

TOPIC TAGS: algorithm, numerical method

ABSTRACT: Algorithms are constructed for the minimization of functionals $\Phi(x)$ of the vector $x(x_1, x_2, \dots, x_p)$ of a Euclidean space E_n with saddle points. The basic algorithm \mathcal{A}_1 is universal for a class S of all functionals $\Phi(x)$ with the

20
19
8

Card 1/2

L 55956-65

ACCESSION NR: AP5010818

\mathbb{R}_k is constructed, and it is proved that, for any polynomial $P(z)$ not having multiple roots, the sequence z_k constructed according to this algorithm will converge to one of the roots of $P(z)$ with any arbitrary initial approximation z_0 . Orig. art. has: 2 formulas.

ASSOCIATION: Fiziko-tehnicheskiy institut nizkikh temperatur Akademii nauk SSSR
(Physico-technical Institute, Academy of Sciences, SSSR)

Card 2/2

L 04205-67 EWT(1)
ACC NR: AP6030006

SOURCE CODE: UR/0020/66/169/005/1026/1029

AUTHOR: Glazman, I. M.; Mitin, V. N.

ORG: Kharkov Polytechnical Institute im. V. I. Lenin (Khar'kovskiy politekhnicheskii institut); Physico-Technical Institute of Low Temperature AN UkrSSR (Fiziko-tekhni-cheskiy institut nizkikh temperatur AN UkrSSR)

TITLE: The tuning out of vibration systems as a problem in convex programming

SOURCE: AN SSSR. Doklady, v. 169, no. 5, 1966, 1026-1029

TOPIC TAGS: linear programming, algorithm, optimal control, resonance damping

ABSTRACT: If Q is a vibration system with m degrees of freedom and m rigidities, and considering this system to be an elastic bar-conductor with rigidities α_k and moments of inertia I_k ($k=1,2,\dots,m$), the problem studied is that of selectively tuning Q out of the resonance danger zone $(0,\beta)$, the interval of squares of possible frequencies from external effects. This problem consists in altering a given number p of its rigidities such that the square λ of the least natural frequency of the system will satisfy the inequality $\lambda \geq \beta$. An algorithm is constructed for the solution of the problem of optimal tuning and also an algorithm for the solution of an analogous problem with supplementary limitations. These algorithms are realized by the use of rational opera-

L 04205-67

ACC NR: AP6030006

tions only, they have an exponential rate of convergence, and they provide for caging in attaining a given degree of accuracy. They are adapted for computer use. The authors thank L. I. Shtenvol' for phasing the problem in terms of the $\lambda \geq 8$ inequality. Presented by Academician A. Yu. Iahlin'skiy on 27 November 1965. Orig. art. has: 14 formulas.

SUB CODE: 12,09/

SUBM DATE: 25Nov65/

ORIG REF: 005/

OTH REF: 001

Card 2/2

GLAZMAN, M.G.; ORLOV, N.G., kandidat meditsinskikh nauk, glavnyy vrach.

Study of stable strains of staphylococci; author's abstract. Zhur. mikro-
biol. epid. i immun. no. 2:53-54 F '53. (MLBA 6:5)

1. Bol'nitsa imeni Baumana.

(Staphylococcus)

USSR/Medicine - Antibiotics

FD-2319

Card 1/1 Pub 146 - 26/55

Author : Glazman, M. G.; Lepikhova, L. F.

Title : Ekmolin as a factor which reinforces the action of penicillin on resistant staphylococcus cultures

Periodical : Zhur. mikro. epid. i immu. No 2, 58, Feb 1955

Abstract : Found that penicillin to which ekmolin has been added has a bacteriostatic effect on staphylococci cultures that are otherwise-resistant to penicillin and that penicillin and ekmolin exert a synergetic bacteriostatic action on staphylococci cultures.

Institution : Hospital imeni Bauman, Moscow

Submitted : March 18, 1954

USSR/Microbiology - Antibiotics and Chemother. Anti-Infect. F 2

Abs Jour : Referat Zhurn - Biol. No 16, 25 Aug 1981, 1981

Author : Glazman, M.G., Belvaeva, L.A.

Title : The Study of Mycelin Activity on Experimental Infection

Orig Pub : Antibiotiki, 1981, 1, No 1, 1-3, 11.

Abstract Of 36 cultures of staphylococci isolated from patients, 35 were sensitive to mycerin, 10 to penicillin, 2 to streptomycin, 31 to biomyxin. Of 27 strains of intestinal bacilli, 21 were sensitive to mycerin, 10 to streptomycin, 10 to biomyxin. Not a single one was penicillin sensitive. Of 94 microbial associations (isolated from the mucus of patients and consisting of grampositive and gramnegative flora), 77 were sensitive to mycerin, 2 to penicillin, 32 to streptomycin, 49 to biomyxin. Of 9 penicillin-resistant strains of proteus vulgaris, 7 were sensitive to mycerin, 4 to streptomycin and 1 to biomyxin. Mycerin was used in concentrations of

Card 1/2

- 28 -

USSR/Microbiology - Antibiosis and Symbiosis. Antibiotics

P-2

Abs Jour : Referat Zhurn - Biol. No 16, 25 Aug 1957. 68470

0.3- 5 /ml. A synergistic action of mycerin with penicillin and streptomycin was established. The most clearly evident synergistic effect was noted in combinations of mycerin with biomycin.

Card 2/2

- 29 -

GLAZMAN, M.G.

~~SECRET~~
Determining plasma coagulation in whole blood. Lab.delo 2 no.5:28
S-O '56. (MIRA 9:11)

1. Iz bol'nitsy imeni Baumana (flavnyy vrach - kandidat meditsinskikh
nauk N.G.Orlov), Moskva.
(BLOOD--COAGULATION)

GLAZMAN, M.G.

Method for determining antibiotic sensitivity of microflora isolated
from the phlegm of the diseased. Lab.delo 2 no.6:17-18 N-D '56.

(MLHA 9:12)

1. Iz bol'nitsy imeni Baumana (glavnyy vrach - kandidat meditsinskikh
nauk M.G.Orlov), Moskva.

(ANTIBIOTICS) (MUCUS) (MICRO-ORGANISMS)

GLAZMAN, N.G.

Detection of yeast cells in sputum during antibiotic therapy. Zhur.
mikrobiol.enid. i immun.,supplement for 1956:41-42 '57 (MIRA 11:1)

1. Iz gorodskoy bol'nitsy No.29 imeni Baumann.
(YEAST) (SPUTUM)

GLAZMAN, M.G.

Comparative effect of neomycin antibiotics with penicillin, streptomycin, and chlortetracycline on some pathogens of suppurative infections [with summary in English]. Antibiotiki 3 no.6:110-111. N-D '58. (MIRA 12:2)

1. Bol'nitsa imeni Bauman (glavnyy vrach - kand. med. nauk N.G. Orlova).

(ANTIBIOTICS)

(SUPPURATION)

(BACTERIA, EFFECT OF DRUGS ON)

BARTSHNIKOVA, O.L.; GLAZMAN, M.G.

Mycerin in clinical practice. Antibiotiki 4 no.3:105-107
My-Je '59. (MIRA 12:9)

1. Gorodskaya bol'nitsa No.29 imeni Baumana (glavnyy vrach
M.G.Orlov).

(ANTIBIOTICS, ther. use,
mycerin (Rus))

GLAZMAN, M.G.

Experimental study of monomycin. Antibiotiki 7 no.5:464-467 My '62,
(MIRA 15:4)

1. Gorodskaya bol'nitsa No.29 imeni Baumana (glavnyy vrach N.G.Orlov).
(ANTIBIOTICS)

GLAZMAN, M.S.

Social and political life at the Institute of Forestry in
1907-1910. Nauch.trudy LTA no.95:115-126 '61. (MIRA 16:2)
(Leningrad--Students) (Leningrad--History)

CHALENKO, N.F.; GLAZMAN, M.Yu.

New developments in the clothing industry enterprises of the Kiev
Economic Council. Leh.prom. no.3:57-59 Je - Ag '62. (MIRA 16:2)

1. Otrasleye konstruktorskoye buyro tresta shveynoy promyshlennosti
Kiyevskogo soveta narodnogo khozyaystva.
(Kiev Economic Region—Clothing industry)

TYAGUNOVA, Z.A.; KUBAREVA, Ya.A.; GLAZMAN, R.A.

Adoption of the continuous neutralization of hydrolyzates at the Krasnodar Hydrolysis Plant. Gidroliz.i sosokhin.prom. 12 no.2:15-17 '59.
(MIRA 12:3)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut gidroliznoy i sul'fitno-spirovoy promyshlennosti (for Tyagunova, Kubareva). 2. Krasnodarskiy gidroliznyy zavod (for Glazman).
(Krasnodar--Hydrolysis)

GLADNEVA, A.N.; GLAZMAN, R.A.; GUREVICH, N.S.; MALINOVSKAYA, Ye.V.

Chemical composition and physical properties of some types of raw
material for hydrolysis. Gidroliz i lesokhim.prom. 12 no.4:
17-20 '59. (MIRA 12:8)

1. Krasnodarskiy gidroliznyy zavod.
(Krasnodar--Hydrolysis)

Coagulation of the alcohol mercuric sulfide. K. Volkov and Yu. Giaman. *Ussr. Khim. Bull. nat. Res. chim.* 1, No. 4, 85-93 (In German) 41, 41 (1945). HgS sol. was prepd. according to Krieger (cf. C. A. 10, 5) to 1 cc. of sol. in 18 cc. of anhyd. EtOH. Add small amounts of electrolyte until a ppt. forms 5 min. after adding the most salts, increase in content hastens coagulation. At small concns., $AlCl_3$ and $FeCl_3$ cause coagulation, but at higher concns. they stabilize the sol. Addn. of benzene increases the mol. polarization of sol. and the sol. becomes unstable. Further study of lyophobic colloids requires the detn. of dipole moments. B. Z. Kamich

A S M - I L A METALLURGICAL LITERATURE CLASSIFICATION

117 AND THE PROJECT		PROCESSES AND PROPERTIES INDEX	
<p>Isomerism of lyophobic colloids. I. The cinnabar alcohol. Yu. M. Glazman and O. M. Baram (Kiev Tech. Inst. Light Ind. J. J. Chem. (U.S.S.R.) 10, 1806-12 (1936) (in Russian)). In a soln. of $\text{Hg}(\text{CN})_2$ in KOH, cooled in an ice-NaCl mixt., prolonged action of H_2S forms a HgS sol of orange-red color (I), different from the usual dark-brown sol (II); the latter does not at any appreciable rate go over into I above -18°, but between -21° and -78° the transformation is complete in 2-4 hrs., the slower the lower the temp. Interruption of the cooling after 5-10 min. satn. of the alcohol of II by H_2S and return to room temp. caused accelerated conversion into I, completed in 10-15 min. Consequently, low temp. is necessary only in the initial stages of the reaction between $\text{Hg}(\text{CN})_2$ and H_2S; the subsequent stage of transition of II into I goes faster at a higher temp. By x-ray diffraction, I was shown to be identical with pure cinnabar, whereas II is known to be structurally metacinnabarite. In MeOH, PrOH, BuOH, and AmOH, transition of II into I takes place in the same temp. range as in KOH. In alic. conig. 32°; H_2O, the threshold temp. for the transition is -25°, with 50% H_2O, -28°; a 10% H_2O-alic. mixt. freezes before the threshold temp. is reached. The conversion does not take place in Me_2CO, the presence of which in an alic. suppresses the II \rightarrow I transition altogether. The detg. factor for the initiation of the conversion appears to be a sufficient concn. of H_2S; in fact, the observed threshold temps. correspond to the same "threshold concn." of H_2S in the various solvents, 1.5-1.6 mole/l.; this concn. cannot be attained in water or in water-rich alic., hence the failure to achieve conversion in these media. This factor does not, however, explain the neg. effect of Me_2CO in which the concn. of H_2S is somewhat higher than in alic. Sol II when formed at a high temp. cannot be converted into I by subseq. cooling; only a sol formed from the very beginning at a low temp. is susceptible of conversion. This reactivity is tentatively ascribed to formation of low-stability HgS \cdot H_2S complexes from which HgI crystallites in the cinnabar lattice, in analogy to a certain extent with the usual artificial prepn. of cinnabar from black HgS by heating with alk. sulfides. The neg. effect of Me_2CO may be due to complex formation with H_2S, producing soln. of H_2S Sol I cannot be obtained at any temp. from soln. of the HgCl_2 or $\text{Hg}(\text{AcO})_2$, probably because of decompn. of the $\text{HgS} \cdot \text{H}_2\text{S}$ complexes by the H^+ ions liberated from H_2S. From this point of view, it is understandable that the conversion is possible only if the H_2S is formed from the un- N. Thon disoord. $\text{Hg}(\text{CN})_2$.</p>		<p>2</p>	
<p>ASB-51A METALLURGICAL LITERATURE CLASSIFICATION</p>			
<p>12000 117000000</p>		<p>12000 117000000</p>	
<p>12000 117000000</p>		<p>12000 117000000</p>	

PROCESSING AND PROPERTIES INDEX									
<div style="position: relative;"> COMMON ELEMENTS COMMON VALUABLE INDEX <div style="position: absolute; left: 0; bottom: 0; width: 100%; height: 100%;"> <div style="position: absolute; top: 10px; left: 10px; font-family: cursive; font-size: large;">CA.</div> <div style="position: absolute; top: 10px; right: 10px; font-size: x-large; font-weight: bold;">2</div> <p style="text-align: center; margin-top: 20px;">Coagulation of lyophobic organosols by electrolytes. I</p> <p>Vin M. Glazman (Technol Inst of Light Industry, Kiev). <i>J. Phys. Chem. U.S.S.R.</i>, 20, 2018, 1946. HgS solns in H₂O, EtOH, COMe, and EtOAc were prepd from Hg(CN)₂ and H₂S. They were coagulated by its different salts. The coagulating power of the cations increased with valency (ratio 1:2:3:4) in the ratio 1:24:545:3000 for water, 1:50:118:222 for EtOH, 1:10:1000 for COMe (valency ratio 1:2:3), and 1:7.5:61 for EtOAc; no quadrivalent ions were used in acetone and ethyl acetate. The <i>elec. cond.</i> of the solns. used is given. II. <i>Ibid.</i>, 209-12. -HgS soln in water and ethanol were coagulated by 13 different Na salts. The largest ratio of the coagulating concn. was 7 in water (for p-O₂NC₆H₄-CO₂Na-NaClO₄) and 2.7 in EtOH (the CH₃CH(OH)-Na-NaBr). It is suggested that fewer anions are adsorbed by HgS in EtOH, with its low dielec. const., than in water.</p> </div> </div>									
ASSOCIATED METALLURGICAL LITERATURE CLASSIFICATION									
FROM SYNONYMS FROM SUBJECTS FROM ANALYSIS FROM METHODS									
SEARCHED BY DATE INDEXED BY DATE ABSTRACTED BY DATE REPRODUCED BY DATE									

27

6

Lyophobic Organic Colloids. (In Russian) L. M. Glazman. *Progress in Chemistry* (USSR), 5, 16, no. 2, 1947, p. 177-188.

Discusses theories concerning the stability of the above systems. Advances tentatively a hypothesis that all lyophobic colloid systems, organic or aqueous, should be considered from the same point of view. 41 ref.

2

Acclimatization effect in the coagulation of hydrophobic colloids by electrolytes. Yu. M. Glazman and H. E. Tartakovskaya. *Kolloid. Zhur.* 11, 200-207 (1949). — A AgI sol (0.01 N + 5% excess of KI) was dialyzed in cellulose bags until its elec. cond. was 8×10^{-4} , and then electrodialed between cellulose membranes until a was 4×10^{-4} ohm⁻¹ cm⁻¹. To 1 ml. of this sol so much coagulating electrolyte (I) was at once added (its final concn. was c_1) that the turbidity of the filtrate after 24 hrs. of coagulation was 1.6 that of H₂O, or I was added 1 drop daily until the filtrate taken 24 hrs. after the last drop had the 1.6 turbidity. In the 2nd case the final concn. c_2 was less than c_1 . From the earlier theory (*Kolloid Zhur.* 9, 241 (1947)) $(c_2 - c_1)/c_1 = (1/k_1) \ln(1 + e^{-k_1 t})$, if $1/n$ is no. of drops used in the 2nd case and k is the const. of coagulation detd. from the exptl. relation between the concn. of I and the time t after which the filtrate had the 1.6 turbidity. For KCl, BaCl₂, and CeCl₃ k was 0.076, 5.12, and 150 l./millimol., and k_1 was 6.2-6.5 for all 3 electrolytes. As k_1 was independent of the nature of I, $(c_2 - c_1)/c_1$ also was identical (0.25 to -0.30) for these salts, and the above equation was confirmed ($1/n$ was 43-46). This proves that the rate of coagulation generally is an exponential function of the concn. of I, although expt. showed that very small amts. of I were less active than expected. In undialyzed AgI sols greater initial amts. of I were inactive, wherefore the above equation yielded greater values of $(c_2 - c_1)/c_1$ than found by expt. J. J. Bikerman

2A

Relation between the adsorption of coagulator ions and their concentration in solution. D. M. Strashenko and Yu. M. Glazman (L. V. Pisarzhevskii Inst. Phys. Chem., Kiev; *Dopovid. Akad. Nauk Ukrain. R.S.R.* 1930, 293; *Izv. Ukrainian Acad. Nauk Ukrain. R.S.R.* 1930, 293). The adsorption of $Hf(NO_3)_3$ (0.01-0.1 M) in coagulation of hydrosols of AgI (10 millimole/l.), HgI₂, and As₂S₃ (8 millimole/l.), was investigated by tagging with radioactive Sr^{90} . With c = concn. of $Sr(NO_3)_2$ corresponding to the coagulation threshold, the limits adsorbed, at the concns. c , $5c$, and $10c$, are: on AgI, 47.0 (micromole/g (4.17%)), 184.7 (3.47), 815.1 (2.90); on HgS, 16.8 (5.10), 46.8 (2.87), 81.3 (2.56); on As₂S₃, 63.9 (12.54), 90.2 (3.56), 118.5 (2.33). The adsorption is practically instantaneous, and the Sr^{90} ions are held firmly by the coagulate. N. Thon

2

CA

Investigation by the tracer method of the coagulation of lyophobic sols by electrolytes. Adsorption of ions of the same sign. Yu. M. Glazman and D. N. Strazhenko (L. V. Pisarzhevskii Inst. Phys. Chem., Kiev). *Doklady Akad. Nauk S.S.S.R.* 75, 411-13(1950). The amt. of neg. ions PO_4^{3-} and SO_4^{2-} adsorbed by the neg. hydrosols of AgI (0.01 M) and HgS and As_2S_3 (0.001 M) was detd. by tagging the anions with radioactive P^{32} and S^{35} , resp. Both at the threshold of coagulation, and in the presence of a twofold excess of the electrolyte the amts. of anion adsorbed were insignificant. PO_4^{3-} was not adsorbed on AgI, and the amts. adsorbed on HgS and As_2S_3 were ~ 0.8 and ~ 0.9 millimole/g., resp.; adsorption of SO_4^{2-} on AgI ~ 0.4 millimole/g., resp.; adsorption of SO_4^{2-} on As_2S_3 and HgS, and ~ 2.1 millimole/g. on As_2S_3 . Adsorption of such small amts. can play no role in coagulation, ion antagonism, etc. Pos. Se^{6+} ions (tagged with radioactive Se^{75}) were adsorbed, at the threshold of coagulation, to the extent of 17.6 millimole/g. on AgI, 16.8 on HgS, and 63.9 on As_2S_3 . N. Thon

1957

Study of the electrostatic regulation of lyophobic sols by
the method of tagged atoms. I. Adsorption of ions having a
charge of the same sign as the particles. Y. M. Glazov,
D. M. Strashenko, and N. E. Tsvetkovskiy. *Chem. Abstr.*
U.S.S.R. 15, 161-71 (1968) (Engl. translation). See C.A.
47: 9107a.

GLAZMAN, Yu.M.

Nature of the stability of lyophobic organosols. Koll.zhur. 15 no.4:225-233 '53. (MLRA 6:8)

1. Tekhnologicheskii institut legkoy promyshlennosti (Kiyev). (Colloids) (CA 47 no.22:11886 '53)

GLAZMAN, Yu.M.

Nature of the "acclimatization" phenomenon during the coagulation of hydrophobic sols by electrolytes. Koll.zhur. 15 no.5:334-346 '53. (MLRa 6:9)

1. Kievskiy tekhnologicheskii inatitut logkoy promyshlennosti.
(Colloids)

GLAZMAN, Yu.M.

Effect of acclimatization during the coagulations of mercuric-sulfide hydrosol by electrolytes. Ukr.khim.zhur. 19 no.5:496-503 '53. (MLBA 8:2)

1. Kiyevskiy tekhnologicheskij institut legkoy promyshlennosti. (Mercuric sulfide)(Coagulation)

GLAZMAN, Yu. M.

USSR/ Chemistry - Colloids

Card 1/2 Pub. 22 - 27/52

Authors : Glazman, Yu. M., and Dykman, I. M.

Title : Coagulation of lyophobic sols under the effect of electrolyte mixtures

Periodical : Dok. AN SSSR 100/2 299-302, Jan 11, 1955

Abstract : It was established that when the process of coagulation of lyophobic sols is not complicated by additional phenomena like adsorption, chemical reaction, etc, the addition to the sol of two electrolytes (differing in valence of the coagulating ions) will be expressed by a plain synergism effect. The phenomena of instances where antagonism may appear are discussed.

Institution : The Technological Institute of Light Industry, Kiev

Presented by : Academician A. N. Frumkin, July 29, 1954

Periodical : Dok. AN SSSR 100/2, 299-302, Jan 11, 1955

Card 2/2 Pub. 22 - 27/52

Abstract : The antagonism was found to be in some way connected with the adsorption reactions which take place during the addition of electrolytes to the sol. The role of the colloidal-dispersion phase in the process of sol coagulation is explained. Seven references: 4 USSR, 2 German and 1 Dutch (1927-1946). Graph.

GLAZMAN, Yu. M.; Zhel'vis, Ye. F.

"On the Nature of the Phenomenon of Assimilation in the Coagulation of Hydrophobic Colloids with Electrolytes. Coagulation of Sol of Prussian Blue (O prirode yavleniya priyemaniya pri kognulyatsii gidrofobnykh kolloidov elektrolitami. Kognulyatsiya zolya Berlinskoy laruri) from the book Trudy of the Third All-Union Conference on Colloid Chemistry, pp. 341-348, Iz. AN SSSR, Moscow, 1966

(Report given at above Conference, Minsk, 21-3 Dec 66)

Authors: Kiev, Technological Institute of Light Industry

GLAZMAN, Yu. M.

Coagulation of lyophobic sols by the action of electrolyte mixtures. Yu. M. Glazman and I. M. Deryagin (Technical Inst. Light Ind., Kiev). Kolloid. Zhur. 18, 13-26 (1956); cf. C.A. 47, 11886; 49, 14739d. --The mutual repulsion of 2 diffuse double layers in a common soln. of a uni-univalent and a bi-valent electrolyte is calcd. by the method of Deryagin and Landau (C.A. 39, 1537). Coagulation occurs when the van der Waals attraction overcomes this repulsion. At all ratios of the concns. of the 2 electrolytes, the mix is a stronger coagulant than is the sum of the components. As this conclusion is not in accord with many observations, the electrostatic repulsion is not the only factor in the coagulation process. 50 references. J. J. Bikerman

PM 5/24/57

AUTHORS: Glazman, Ya. M., Dymchenko, I. M., Strel'tsova, T. A., 20-117-5-29/54

TITLE: The Antagonism of Ions in the Coagulation of Lyophobic Sols by electrolytes (Ob antagonizme ionov pri kognulyatsii lyofobnykh zolei elektrolitami).

PERIODICAL: Doklady AN SSSR, 1957, Vol. 117, Nr 5, pp. 829-832 (USSR)

ABSTRACT: In a preliminary paper of the authors (reference 1) it was shown, that on the coagulation of lyophobic sols by a mixture of two symmetric electrolytes of the type 1 - 1 + 2 - 2 the synergism effect must be taken into consideration almost in the whole range of their concentrations. At the same time the supposition was pronounced, that in the case of a differing composition of the electrolytes the computations may furnish quite different results. The present investigation is destined to examine the correctness of this assumption. The authors conducted computations analogous to the ones mentioned in the preliminary paper, especially for the case of the coagulation of a sol by a mixture of electrolytes of the type 12 - 2 + 2 - 2. At first a differential equation for the electrolytic potential in an arbitrary point of the solution is given. This potential Ψ is then, for the sake of convenience, replaced by a dimensionless potential. Besides, the sol is here supposed to be strongly charged. The whole range of the possible concentrations n_1 and n_2 is of identical interest for the problem

Card 1/2

The Antagonism of Ions in the Coagulation of Lyophobic Sols by Electrolytes. 20-117-5 -29/54

Investigated here. The parameter n_1 is here considered to be infinitely small, and it is sufficient to break off the series expansion with respect to n_1 after the first term. Then formulae corresponding to this approximation are written down. The course of the computation is followed step by step and shows the subsequent results; The rules of the coagulation of lyophobic sols by mixture of electrolytes of the type 1 - 1 +2 -2 and 12 -2 +2 - 2 are opposed to each other in a qualitative sense. In the first case the synergism is substantiated theoretically and in the second case the theory leads to a sharply pronounced antagonism. In the coagulation of lyophobic colloids there must be distinguished two types of antagonisms: 1) An antagonism connected with the competition for the adsorption places on the surface of the colloidal particles. 2) An antagonism caused by the electrostatic interaction of the ions in the volume of the solution and in the electric field of the colloidal particles. There are 3 references, 2 of which are Slavic.

PRESENTED: June 10, 1957, by P. A. Rasinder, Academician

SUBMITTED: June 7, 1957

Card 2/2

GLAZMAN, Yu. V.

AUTHORS: Glazman, Yu. V., Strazhesko, D. N., Lisikalova, V. A. 73-1-21/43

TITLE: Investigation of the Coagulation of Lyophobic Sols Through Electrolytes by Means of the Method of Marked Atoms (Issledovaniye koagulyatsii liofobnykh zolei elektrolitami metodom mechenykh atomov).
II. Adsorption of Cations by Positively Charged Colloidal Particles (II. Adsorbtsiya kationov polozhitel'no naryazhennymi kolloidnymi chastitsami).

PERIODICAL: Zhurnal Neorganicheskoy Khimii, 1958, Vol. 3, Nr 1, pp. 115-118 (USSR).

ABSTRACT: The aforesaid coagulation is always accompanied by phenomena of adsorption. The question of the importance of the latter is still of topical interest for the theory of the stability of lyophobic colloids. It is, in particular, not clear whether an adsorption of indifferent ions which have a charge of the same sign as the colloidal particles, takes place and which rôle this adsorption plays in the mechanism of coagulation. The authors state in a previous report that the adsorption of indifferent anions was very small (up to 1 to 2 micro mol per 1 g disperse phase) with the coagulation of negative brine by electrolytes. It was interesting to clarify, in which way the cations will behave in coagulation in this respect, as mentioned in the sub-title. The present

Card 1/1

Investigation of the Coagulation of Lyophobic Sols Through
Electrolytes by Means of the Method of Marked Atoms.

72-1-21/43

report is devoted to this problem. A voluminous experimental part follows. Positively charged brine of ferric hydroxide and silver iodide served for the investigation. The dialysis was carried out in little bags of cellophane with frequent change of the water in the exterior vessel, for 14 days. The chlorides of potassium, rubidium, caesium, calcium, which were correspondingly marked with K^{42} , Rb^{86} , Cs^{134} , Ca^{45} served for the coagulation, as well as nitrates of silver and lanthanum (marked with Ag^{110} and La^{140}) served for the coagulation. The coagulating concentrations were visually determined (like in reference 1). The minimum concentration which was sufficient for achieving the complete separation of the disperse sol phase from the medium of dispersion, was denoted as threshold of coagulation ($C_{K.S.} = C_{p.k.}$). The usual (commercial) radio isotopes lead, in spite of quite small quantities of contaminations, to somewhat unexpected results which substantially differed from those given in the previous report (reference 1): The adsorption of cations was quite important in several cases. The isotopes were therefore - in addition - still especially purified. The results obtained with such purified radioactive preparations are given in table 1 and 2. They show that the adsorption of cations with the coagulation of positively charged brine

Card 2/4

Investigation of the Coagulation of Lyophobic Sols Through Electro- 78-1-21/43
lytes by Means of the Method of Marked Atoms.

of ferric hydroxide and silver iodide is extremely small and that it increases very slightly when a surplus of the coagulating electrolyte is added to the colloidal solutions. Rubidium-, caesium-, and calcium-ions are practically not adsorbed at all. The adsorption of potassium-ions seems to be somewhat higher at first sight, but in reality any radiochemical admixture was adsorbed from which the authors could apparently not completely liberate the solution. The ion-adsorption of lanthanum and silver was somewhat higher, apparently due to their specific adsorption power which is caused by structural peculiarities of their electron shells (reference 8). Consequently, the results of the present report prove, as well as those of the previous paper (reference 1) that the adsorption of ions of the same sign, especially with the coagulation of positively charged brine, is extremely small. Immediate radiometric measurements of the adsorption of the opposed ions are required, however, for a final judgement of the small adsorption power of ions of the same sign and their rôle in the mechanism of various phenomena of coagulation. There are 2 tables, and 13 references, 8 of which are Slavic.

Card 3/4

Investigation of the Coagulation of Lyophobic Sols Through
Electrolytes by Means of the Method of Marked Atoms.

7-1-21/13

ASSOCIATION: Technological Institute for Light Industry (Tekhnologicheskii Institut Legkoy Promyshlennosti).
Medical Institute im. A. A. Bogomolets, Kiev (Meditsinskiy Institut im. A. A. Bogomol'tsa, Kiev)

SUBMITTED: May 18, 1957.

AVAILABLE: Library of Congress.

Card 4/4

69-58-2 -4/23

AUTHORS: Glazman, Yu.M.; Dykman, I.M.; Strel'tsova, Ye.A.

TITLE: The Coagulation of Lyophobic Sols by the Action of Electrolyte Mixtures. Communication 2. (O koagulyatsii lyofobnykh soley pri deystvii smesey elektrolitov. Soobshcheniye 2)

PERIODICAL: Kolloidnyy zhurnal, 1958, Vol XX, Nr 2, pp 149-158 (USSR)

ABSTRACT: During coagulation of lyophobic sols, caused by the mixture of two symmetrical electrolytes of the 1-1+2-2 type, a synergistic effect takes place within their concentration range. The action of these two electrolytes is accompanied by two contradicting factors. Synergism is caused by the addition of the electrolyte and is connected with the compression of a diffused ion atmosphere. A tendency toward antagonism is caused by the screening of the anions of the electrical field. Electrolytes of the 1₂-2+2-2 type have also been considered. The principal difference of the two types consists in the fact that the side ion of the first of the two electrolytes is bivalent. The theoretical calculation indicates antagonism, whereas in the 1-1+2-2 type it indicates synergism. There are two types of antagonism during the coagulation of lyophobic colloids: 1) the antagonism between

Card 1/2

69-58-2 -4/23

**The Coagulation of Lyophobic Sols by the Action of Electrolyte Mixtures.
Communication 2**

coagulating ions associated with competition for the sites of adsorption on the surface of the colloid particles: 2) the antagonism due to the mutual electrostatic interaction of the ions in the bulk of the solution and in the electrical field of the colloid particles.

There are 8 references, 6 of which are Soviet, 1 French and 1 German.

ASSOCIATION: Kiyevskiy tekhnologicheskii institut legkoy promyshlennosti
(Kiev Technological Institute of Light Industry)

SUBMITTED: February 23, 1957

1. Chemical compounds--Coagulation 2. Electrolytic compounds
--Applications

Card 2/2

GLAZMAN, Yu. M.

"The first mathematical theory of ion antagonism."

report presented at the Fourth All-Union Conference on Colloidal Chemistry,
Tbilisi, Georgian SSR, 12-16 May 1958 (Koll zhur, 20,5, p.877-9, '58, Traubman, A.B)

9.12.88, Vol. 1., Doc Chem Sci — (disc) "Study of the stability
and solvation of lyophilic solids in aqueous electrolytes." Div,
195 . 23 pp with append (Inst of General and Inorganic Chemistry
of the Acad Sci USSR). 110 copies. List of author's works, p. 1-23.
(13,27-1, 10)

DOI/47-1-1-5/25

5(4)

AUTHORS: Glazman, Yu.M., Strazhenko, D.M., Zhel'vis, Ye.F.,
Chervyatsova, L.L.

TITLE: Changes in the Adsorption of Potential-Determining
Ions During Coagulation of Lyophobic Sols by In-
different Electrolytes

PERIODICAL: Kolloidnyy zhurnal, 1979, Vol XXI, Nr 5, pp 265-271
(USSR)

ABSTRACT: The present investigation concerns the role of the
potential-determining ions during the coagulation
process of lyophobic sols, caused by indifferent
electrolytes with coagulating ions of different
valency. Objects of the investigation are the ac-
tivity of the investigated sol: As_2S_3 (negatively charged)
and $Fe(OH)_3$ (positively charged). Comparing the
magnitudes of the activities of the intercellular
liquids of the investigated sols with the activities
of the corresponding solutions after coagulation,

Card 1/3

Changes in the Adsorption of Potential-Determining Ions During
 Congulation of Lyophobic Sols by Indifferent Electrolytes

SOV/15-11-1-3/15

A marked additional adsorption of potential-determining ions could be stated in other sol. The desorption of iron ions, which could be observed during the congulation of the Fe(OH)₃ sol, was due to secondary factors. Congulation of lyophobic sol by indifferent electrolytes, therefore, affects not only the external but also the internal surface of the colloidal particle double layer. The mechanism observed thereby cannot be explained from the standpoint of a purely electrostatic compression of the double layer. There is a quantitative discrepancy between this conception and the obtained data. The authors conclude by recommending the further study of the congulation theory, which is to consider the quantitative effect of electrolytes on the surface potential of colloidal particles. Towards the end of the article, the authors mention the Soviet scientists V.A. Margolin and A.I. Rabinovich in connection with certain effects produced by poten-

Card 2/3

SOV/69-21-3-3/25

Changes in the Adsorption of Potential-Determining Ions During
Coagulation of Lyophobic Sols by Indifferent Electrolytes

tial-determining ions during the coagulation process.
There are 3 tables and 50 references, 24 of which are
Soviet, 13 German, 10 English and 3 French.

ASSOCIATION: Tekhnologicheskii institut legkoy promyshlennosti
(Technological Institute of Light Industry)
Institut fizicheskoy khimii AN USSR im. L.V. Pissar-
zhevskogo, Kiev (Institute of Physical Chemistry
of the AS of the USSR imeni L.V. Pissarzhevskiy,
Kiev)

SUBMITTED: 26 February 1958

Card 3/3

GLAZMAN, Ya. M. and STRAZHEK, D. N.

"The Significance of Adsorption Phenomena in the Mechanism of Localization of
Lyophobic Colloids by Electrolytes."

report presented at the Section on Colloid Chemistry, VII. Mendeleev Conference on
General and Applied Chemistry, Moscow, 10-13 March 1967.
(Koll. Zhur. v. 31, No. 6, pp. 1071-81)

С. Л. ДИМИТОВ, В. И. ДИМИТОВ

PHASE I BOOK EXPLORATION

07/01/90

Konferentsiya po poverkhnostnykh silam. Moscow, 1969.

Issledovaniya v oblasti poverkhnostnykh sil; sbornik dokladov na konferentsii po poverkhnostnykh silam, april' 1969 g. (Studies in the Field of Surface Forces; Collection of Reports of the Conference on Surface Forces, Held in April 1969) Moscow, Izdatel'stvo AN SSSR, 1969. 231 p. Errata printed on the inside of back cover. 2500 copies printed.

Sponsoring Agency: Institut Fizicheskoy Khimii Akademii nauk SSSR.

Revs. Ed.: M. V. Deryagina, Corresponding Member, Academy of Sciences USSR; Editorial Board: M. M. Zeldavayev, V. A. Krut'ko, N. N. Kuznetsov, S. V. Merzlin, P. S. Prokhorov, M. V. Salayev and G. I. Pains; Ed. of Publishing House: A. L. Bankvitsker; Tech. Ed.: Yu. V. Rykina.

PURPOSE: This book is intended for physical chemists.

Card 1/8

Studies in the Field of Surface Forces (Cont.)

201/0555

CONTENTS: This is a collection of 25 articles in physical chemistry on problems of surface phenomena investigated at or in association with the Laboratory of Surface Phenomena of the Institute of Chemical Chemistry of the Academy of Sciences USSR. The first article provides a detailed chronological account of the Laboratory's work from the day of its establishment in 1935 to the present time. The remaining articles discuss general surface forces, problems of polymer adhesion, surface forces in thin liquid layers, surface phenomena in dispersed systems, and surface forces in aerosols. Names of scientists who have been or are now associated with the Laboratory of Surface Phenomena are listed with references to their past and present associations. Each article is accompanied by references.

TABLE OF CONTENTS:

Zakhar'yeva, N. N. Twenty-Five Years of the Laboratory of Surface Phenomena of the USSR (Institute of Physical Chemistry of the Academy of Sciences USSR)

3

Card 2/6